

## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <a href="http://about.jstor.org/participate-jstor/individuals/early-journal-content">http://about.jstor.org/participate-jstor/individuals/early-journal-content</a>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

finds that there is little reason to suppose that they are of any service in protecting the flowers from ants.—C. R. B.

Centrosomes in Marchantia.<sup>36</sup>—After a study of spermatogenesis in *Marchantia polymorphia*, Schaffner concludes that Ikeno's account is correct and that centrosomes are present, both while the nucleus is at rest and while it is undergoing mitosis. His figures are practically the same as Ikeno's. Miyake's failure to find centrosomes he attributes to differences in technic.—Charles J. Chamberlain.

Thermotropism.—POHL<sup>37</sup> describes observations and experiments that he has made upon the cultivated flax, which show its great sensitiveness to radiant heat, the young shoots directing themselves toward the source. Experiments also show that heat is the dominant factor in inclination of the shoots, which is often ascribed to light.—C. R. B.

Discomycetes.—MISS BACHMAN has published<sup>38</sup> a descriptive catalogue of the Discomycetes within five miles of Oxford, O. Keys to genera and species are provided, and there are illustrations of ten of the sixty-odd species, a goodly number of which are now for the first time reported from southwestern Ohio.—C. R. B.

Light and respiration.—Löwschin reports<sup>39</sup> that when he excluded the effects of actinic warming he was unable, in the course of twenty-two experiments upon the respiration of certain fungi (Aspergillus, Penicillium, Oidium, and Cladosporium), to detect any regular acceleration of it by light.—C. R. B.

<sup>&</sup>lt;sup>36</sup> Schaffner, John H., The centrosomes of *Marchantia polymorphia*. Ohio Naturalist **9:**<sub>3</sub>83–388. *pl.* 21. 1908.

<sup>37</sup> Pohl, J., Der Thermotropismus der Leinpflanze. Beih. Bot. Centralbl. 24:111-131. figs. 6. 1908.

<sup>&</sup>lt;sup>38</sup> BACHMAN, FREDA M., Discomycetes in the vicinity of Oxford, Ohio. Proc. Ohio State Acad. Sci. **5**:19–70. *pls.* 4. 1909.

<sup>&</sup>lt;sup>39</sup> Löwschin, A., Zur Frage über den Einfluss des Lichtes auf die Atmung der niederen Pilze. Beih. Bot. Centralbl. 23:54–64. 1908.